CHEST

Official publication of the American C ollege of Chest Physicians



Continuing Medical Education Effect on Practice Performance

Dave Davis and Robert Galbraith

Chest 2009;135;42S-48S DOI 10.1378/chest.08-2517

The online version of this article, along with updated information and services can be found online on the World Wide Web at: http://www.chestjournal.org/content/135/3 suppl/42S.full.html

CHEST is the official journal of the American College of Chest Physicians. It has been published monthly since 1935. Copyright 2007 by the American College of Chest Physicians, 3300 Dundee Road, Northbrook IL 60062. All rights reserved. No part of this article or PDF may be reproduced or distributed without the prior written permission of the copyright holder.

(http://www.chestjournal.org/site/misc/reprints.xhtml) ISSN:0012-3692



CME: ACCP EVIDENCE-BASED EDUCATIONAL GUIDELINES

Continuing Medical Education Effect on Practice Performance*

Effectiveness of Continuing Medical Education: American College of Chest Physicians Evidence-Based Educational Guidelines

Dave Davis, MD; and Robert Galbraith, MD, MBA

Background: There has been sizable debate and widespread skepticism about the effect of continuing medical education (CME) on the performance of physicians in the practice setting. This portion of the review was undertaken to examine that effect.

Methods: The guideline panel used data from a comprehensive review of the effectiveness of CME developed by The Johns Hopkins Evidence-based Practice Center, focusing on the effect of CME on clinical performance.

Results: The review found 105 studies, which evaluated the impact of CME on short- and long-term physician practice performance. Nearly 60% met objectives relative to changing clinical performance in prescribing; screening; counseling about smoking cessation, diet, and sexual practices; guideline adherence; and other topics. Single live and multiple media appeared to be generally positive in their effect, print media much less so. Multiple educational techniques were more successful at changing provider performance than single techniques. The amount or frequency of exposure to CME activities appeared to have little effect on behavior change.

Conclusions: Overall, CME, especially using live or multiple media and multiple educational techniques, is generally effective in changing physician performance. More research, however, is needed that focuses on the specific types of media and educational techniques that lead to the greatest improvements in performance.

(CHEST 2009; 135:42S-48S)

Key words: clinical performance; continuing medical education; guideline implementation; impact of continuing medical education; knowledge translation; performance change; physician behavior; physician performance

Abbreviations: CME = continuing medical education; EPC = evidence-based practice center

SUMMARY OF RECOMMENDATIONS

1. General: We recommend that CME interventions be used to improve physician practice performance (Grade 1C).

*From the Association of American Medical Colleges (Dr. Davis), Washington, DC; and the Center for Innovation (Dr. Galbraith), National Board of Medical Examiners, Philadelphia, PA. Manuscript received October 20, 2008; revision accepted December 10, 2008.

Reproduction of this article is prohibited without written permission from the American College of Chest Physicians (www.chestjournal.org/misc/reprints.shtml).

Correspondence to: Dave Davis, MD, Continuing Health Care Education and Improvement, Association of American Medical Colleges, Washington, DC 20037; e-mail: ddavis@ aamc.org

DOI: 10.1378/chest.08-2517

2. Instructional media:

- a. We recommend that both single live and multiple media be used to maintain or improve physician practice performance (Grade 1C).
- b. We recommend that print media should not be used alone to improve physician practice performance (Grade 1C).
- 3. Instructional techniques: We recommend that multiple instructional techniques be used to improve or maintain physician practice performance (Grade 1C).
- 4. Frequency of exposure: We suggest that CME activities that include multiple exposures, as opposed to a single exposure, be

used to improve physician practice performance (Grade 2C).

There has been considerable debate and widespread skepticism about the effect of continuing medical education (CME) on the performance of physicians in the practice setting. The relationship between education and performance is of great importance to health service managers, quality improvement coordinators, guideline implementers, licensing bodies, policymakers, practitioners, and others. In addition, the suggestion that CME can and should lead to change in clinical performance is an increasingly important principle of the accreditation process used by the Accreditation Council for Continuing Medical Education and to systems of relicensure, recertification, and many other stakeholders in the US healthcare system.

Implications to the physician-learner from the literature review in this article suggest similar findings to those in the preceding two articles. The primary focus here, however, is on physician practice performance as opposed to knowledge acquisition or application. Here again, the guideline panel suggests that live, face-to-face educational activities are effective, especially when combined with multiple exposures to the information following the live educational activity. To the physician-teacher and those who depend on findings from CME activities, the type and amount of multiple exposures are still speculative at best. This article also emphasizes the need for additional studies to be conducted by physician-teachers to compare the number of exposures following a live education activity that are most effective in improving physician performance.

MATERIALS AND METHODS

The guideline panel reviewed the evidence tables and comprehensive review of the effectiveness of CME developed by The Johns Hopkins Evidence-based Practice Center (EPC), which is detailed in the methods article. 2a For this article, we focused on the effect of CME on physician practice performance. These effects were measured in either the short term (< 30 days after an intervention) or the long term (\geq 30 days postintervention). The panel noted several limitations in the methods of the EPC review, which are detailed in the "Science of Continuing Medical Education" article of this supplement and at the conclusion of this article.

RESULTS

Overall Effects of CME on Clinical Performance

The review identified 105 studies, which evaluated the impact of CME on short- and long-term physician practice performance. The majority (61 studies, 58%) met practice objectives. A wide mix of objectives was studied, including prescribing; screening; counseling about smoking cessation, diet, and sexual practices; guideline adherence; and others. Fifty studies^{3–51} were successful in meeting many objectives and demonstrated long-term effectiveness. Among these, evaluation duration ranged from 30 days to 6 months (17 studies) to 1 year or longer (30 studies).

The majority of studies reported positive outcomes, but slightly less than 30% did not. Of these, $24^{7,18,27,29,33,34,52-69}$ analyzed their outcomes for > 30 days, $2^{70,71}$ had shorter evaluation periods; and 3^{72-74} did not report their duration of evaluation. Nine studies^{58,59,75-81} showed mixed results, whereas one¹⁸ was unclear about whether it met objectives. Fourteen studies⁸²⁻⁹⁵ lacked a control group; thus, no reliable measurement of effect could be made.

Recommendation

1. We recommend that CME interventions be used to improve physician practice performance (Grade 1C).

Effects of Instructional Media on Clinical Performance

We posed several questions in this process. Is there a difference in the effect of the use of single vs multiple media? What is the effect of the use of a specific media in CME on practice performance? Are some media better or worse than others in changing performance? Twenty studies used single live media of which one half met their performance objectives. Nine of these studies^{3,6,23,35,39,45,51,107,108} reported long-term performance outcomes, suggesting that the use of single live media methods can generate long-term effects on practice performance objectives; one 97 study assessed outcome < 30 days and met its objectives. Three studies^{75,79,80} using single live events reported mixed results, and three^{60,70,72} did not change practice behavior. Finally, four^{85,86,90,91} studies lacked a control group.

Single print media, particularly comprehensive, unsolicited materials, are not effective in the short-or long-term achievement of practice performance objectives. Nine studies examined the impact of single print media. One⁹⁸ met objectives but did not report evaluation duration, whereas four^{52,55,63,66} did not meet objectives over the long term. Four further studies did not meet objectives, with two^{71,74} not specifying an evaluation period and two^{93,95} lacking a control group. Data available on the use of other single media are scant. One study⁶⁸ using an Internet medium reported a long-term evaluation, but it was

unclear whether it met objectives. Three studies either did not meet objectives,⁷³ lacked a control group,^{82,87} or did not report the methods used.⁸⁷

Of 57 studies using multiple media, the majority studies) met their objective. Of these, 314,7-10,12-20,22,24,31-34,36-38,42-44,46,48-50,99 included long-term evaluations, suggesting that multiple media methods can have a favorable long-term effect on practice behaviors. Nine studies^{96,100–107} met objectives using multiple media but did not specify the timing of evaluation. In contrast, 4 studies^{59,76–78} showed mixed results over the long $14^{7,18,33,34,53,54,56,57,59,61,64,65,67,69}\\$ achieve their objectives, and 383,88,92 did not include a control group. Overall, most studies suggested that CME activities that use multiple media have both a short- and a long-term effect on practice behavior objectives.

Fifteen studies provided an opportunity for direct comparisons of single and multiple media methods. Of these, $10^{5,21,25,27-30,41,46,47}$ met 11 objectives, and all were evaluated over the long term. These studies suggest that both single and multiple media methods have a positive short- and long-term effect on practice behavior objectives, although many were methodologically flawed and did not permit direct comparisons or a recommendation.

Recommendations

2. Instructional media:

- a. We recommend that both single live and multiple media be used to maintain or improve physician practice performance (Grade 1C).
- b. We recommend that print media should not be used alone to improve physician practice performance (Grade 1C).

Effects of Educational Techniques on Clinical Performance

The EPC review also focused on the use and effects of specific educational techniques employed within the context of a single medium, such as a live activity. What is the effect of specific CME educational techniques on practice performance? Is there a difference in outcome when one technique is used compared to instances in which multiple techniques are used? The techniques studied included academic detailing, audience response systems, case-based learning, clinical experiences, demonstrations, discussion groups, feedback, lectures, mentoring or precepting programs, point-of-care techniques, problem-based learning, team-based learning, programmed learning, readings, role play, simulations with standardized patients, and writing.

The Effect of Single Educational Techniques

Eleven studies evaluated the impact of a single technique on practice performance. Three of these studies 14,97,98 displayed some evidence of positive changes, although only one 14 met objectives in the long term. In contrast, 4 studies 52,64,71,73 did not meet objectives using a single technique, 1^{80} yielded mixed results, and $3^{85,91,95}$ lacked a control group. In summary, these studies suggest that a short- or long-term positive effect on practice behavior objectives is not clearly achieved through a single technique.

The Effect of Multiple Educational Techniques

A total of 76 studies with 98 objectives evaluated the short- and long-term impact of multiple techniques on practice behavior objectives. Of this number, more than half (47 studies) achieved their objectives. Of these 47, 39^{4-10,12,13,15-21,23,24,26,31-34,36-39,42-51,108} measured their effects over the long term. A further eight^{96,100,101,103-107} used multiple techniques but did not report the timing of evaluation. In contrast, 16 studies^{7,18,33,53-57,59-61,65-69} evaluated long-term outcomes but did not meet objectives, and 3^{70,72,74} that did not specify evaluation timing failed to meet objectives. Fifteen studies^{59,75-79,83,84,87,89,90,92,93,109,110} did not permit interpretation; they were methodologically flawed, did not describe evaluation timing, lacked a control group, or demonstrated mixed results.

Comparison of Single and Multiple Techniques

Eighteen studies^{3,25,27–30,35,40,41,58,62,63,81,82,86,94,99,102} compared the use of single and multiple educational techniques in CME. Ten^{3,25,27-30,35,40,41,99} of these met their objectives over the long term and suggested that multiple techniques may have a more advantageous short- and long-term effect on practice behavior objectives. Two studies^{58,81} reported mixed results, whereas an additional four^{29,58,62,63} either lacked a control group or did not meet objectives. Two^{82,94} studies were methodologically flawed. In summary, the evidence from these studies in which direct comparisons were possible indicates that CME activities that use multiple educational techniques may have a greater overall positive short-term effect, long-term effect, or both on practice behavior objectives than those that use only a single technique.

Recommendation

3. We recommend that multiple instructional techniques be used to improve or maintain physician practice performance (Grade 1C).

The Effect of Different Levels of Exposure to CME Single, One-Time Exposures

Thirty-seven studies evaluated the impact of single exposure to the CME activity on practice performance. Of these, just under half (18 studies) $^{4,5,9,21,23,25-27,38,39,41,43,45,46,51,100,101,108}$ demonstrated a positive effect. In contrast, 8 studies $^{27,52,61,62,64,69-71}$ did not meet their objectives, $5^{85,86,88,90,91}$ lacked a control group, and $6^{75,76,78-81}$ showed mixed results. Therefore, a single CME exposure may have a positive short- and long-term effect on practice behavior objectives.

Multiple Exposures

The impact of multiple exposures to the CME activity was evaluated in 55 studies encompassing 72 objectives. In this set of studies, almost two thirds (35 studies) met their objectives. Of these, 30 studies^{3,6,8,10,12–20,22,24,28,31–36,40,42,44,47–50,99} continued to meet their objectives over the long term. A further five studies^{96,98,103,104,107} met objectives but did not report evaluation timing, and two studies^{58,59} displayed mixed results. In contrast, one third (18 studies) did not achieve an impact on performance. Two of these studies^{72,73} did not report evaluation duration, whereas 16^{18,33,34,53,55–60,63,65–68,111} with long-term evaluation periods did not realize their objectives. Six studies^{82,85,86,88,90,91} lacked a control group.

Comparison of Single and Multiple Exposures to CME on Performance

Eight studies^{7,29,30,77,83,92,94,105} offered direct comparisons between single and multiple exposures; however, they lacked control groups, showed mixed results, or both. Thus, no strong conclusions are possible from this portion of the EPC review. Overall, the data suggest that multiple exposures to CME activities may have more positive effects than single exposures on practice performance in both the short and the long term.

Recommendation

4. Frequency of exposure: We suggest that CME activities that include multiple exposures, as opposed to a single exposure, be used to improve physician practice performance (Grade 2C).

DISCUSSION

It has long been accepted that CME is an important educational intervention for the improvement of practice performance in health care, but the evidence from previous systematic reviews and other data offers less than strong support to this belief.^{1,2,112} Direct comparisons between the EPC review and these studies is made problematic by the EPC's inclusion of self-reported performance outcomes in some studies. This use of self-report with its potential for bias is in contrast to other systematic reviews, which limited the outcomes to objective measures of performance change only.

Nevertheless, the guideline panel accepted the robustness of the EPC report that suggests a more positive interpretation of the effect of CME, noting several themes in the context of CME's role in improving or maintaining practice performance. First, in the broad construct of CME as a live, face-to-face activity, evidence indicates that this form of education may be effective, especially in settings that permit multiple exposures. Second, the review shows improved results with the use of multiple media and multiple educational techniques; this finding is entirely consistent with principles of adult learning theory. 113,114 Third, it is clear that much useful evidence is still missing in terms of related research questions and the more routine inclusion of measures of practice performance in CME activities. Among the missing data is sufficient comparative evidence on which to derive conclusions about such issues as the type and nature of media, the type and nature of educational techniques (eg, case-based learning, interactivity, sequencing of multiple activities), the confounders of learner motivation and setting, and the degree of change required. These questions and a further discussion about possible limitations to the review are framed in the mandate for establishing a robust research agenda, as discussed in "The Science of Continuing Medical Education" article in this supplement.^{2b}

CONFLICT OF INTEREST DISCLOSURES

Dr. Davis has received grants from the university of Toronto for the redevelopment of a database and \$100,00 in grants from the Ministry of Health, Ontario, Canada.

Dr. Galbraith has no conflicts of interest to disclose.

REFERENCES

- 1 Davis DA, Thomson MA, Oxman AD, et al. Changing physician performance: a systematic review of the effect of continuing medical education strategies. JAMA 1995; 274:700-705
- 2 Grimshaw JM, Thomas RE, MacLennan G, et al. Effectiveness and efficiency of guideline dissemination and implementation strategies. Health Technol Assess 2004; 8:iii-iv:1-72
- 2a Marinopoulos SS, Baumann MH. Methods and definitions of terms: effectiveness of continuing medical education: American College of Chest Physicians evidence-based educational guidelines. Chest 2009; 135(suppl):17S-28S

- 2b Davis D, Bordage G, Moores LK. The science of continuing medical education: terms, tools, and gaps: effectiveness of continuing medical education: American College of Chest Physicians evidence-based educational guidelines. Chest 2009; 135(suppl):8S-16S
- 3 Adams A, Ockene JK, Wheller EV, et al. Alcohol counseling: physicians will do it. J Gen Intern Med 1998; 13:692–698
- 4 Anderson JF, McEwan KL, Hrudey WP. Effectiveness of notification and group education in modifying prescribing of regulated analgesics. Can Med Assoc J 1996; 154:31–39
- 5 Beaulieu M, Choquette D, Rahme É, et al. CURATA: a patient health management program for the treatment of osteoarthritis in Quebec; an integrated approach to improving the appropriate utilization of anti-inflammatory/analgesic medications. Am J Manag Care 2004; 10:569–575
- 6 Brown JB, Boles M, Mullooly JP, et al. Effect of clinician communication skills training on patient satisfaction: a randomized, controlled trial. Ann Intern Med 1999; 131:822–829
- 7 Browner WS, Baron RB, Solkowitz S, et al. Physician management of hypercholesterolemia: a randomized trial of continuing medical education. West J Med 1994; 161:572–578
- 8 Bunting PS, Van Walraven C. Effect of a controlled feed-back intervention on laboratory test ordering by community physicians. Clin Chem 2004; 50:321–326
- 9 Chassin MR, McCue SM. A randomized trial of medical quality assurance: improving physicians' use of pelvimetry. JAMA 1986; 256:1012–1016
- 10 Chodosh J, Berry E, Lee M, et al. Effect of a dementia care management intervention on primary care provider knowledge, attitudes, and perceptions of quality of care. J Am Geriatr Soc 2006; 54:311–317
- 11 Chung S, Mandl KD, Shannon M, et al. Efficacy of an educational Web site for educating physicians about bioterrorism. Acad Emerg Med 2004; 11:143–148
- 12 Clark NM, Gong M, Schork MA, et al. Impact of education for physicians on patient outcomes. Pediatrics 1998; 101:831–836
- 13 Clark NM, Gong M, Schork MA, et al. Long-term effects of asthma education for physicians on patient satisfaction and use of health services. Eur Respir J 2000; 16:15–21
- 14 Cohn BA, Wingard DL, Patterson RC, et al. The National DES Education Program: effectiveness of the California Health Provider Intervention. J Cancer Educ 2002; 17:40–45
- 15 Cummings SR, Coates TJ, Richard RJ, et al. Training physicians in counseling about smoking cessation: a randomized trial of the "Quit for Life" program. Ann Intern Med 1989; 110:640-647
- 16 Gerstein HC, Reddy SS, Dawson KG, et al. A controlled evaluation of a national continuing medical education programme designed to improve family physicians' implementation of diabetes-specific clinical practice guidelines. Diabet Med 1999; 16:964–969
- 17 Cummings SR, Richard RJ, Duncan CL, et al. Training physicians about smoking cessation: a controlled trial in private practice. J Gen Intern Med 1989; 4:482–489
- 18 Davis RS, Bukstein DA, Luskin AT, et al. Changing physician prescribing patterns through problem-based learning: an interactive, teleconference case-based education program and review of problem-based learning. Ann Allergy Asthma Immunol 2004; 93:237–242
- 19 Dietrich AJ, Olson AL, Sox CH, et al. Sun protection counseling for children: primary care practice patterns and effect of an intervention on clinicians. Arch Fam Med 2000; 9:155–159
- 20 Gifford DR, Mittman BS, Fink A, et al. Can a specialty society educate its members to think differently about clinical decisions? Results of a randomized trial. J Gen Intern Med 1996; 11:664–672

- 21 Gonzales R, Steiner JF, Lum A, et al. Decreasing antibiotic use in ambulatory practice: impact of a multidimensional intervention on the treatment of uncomplicated acute bronchitis in adults. JAMA 1999; 281:1512–1519
- 22 Harris SB, Leiter LA, Webster-Bogaert S, et al. Teleconferenced educational detailing: diabetes education for primary care physicians. J Contin Educ Health Prof 2005; 25:87–97
- 23 Herbert CP, Wright JM, Maclure M, et al. Better Prescribing Project: a randomized controlled trial of the impact of case-based educational modules and personal prescribing feedback on prescribing for hypertension in primary care. Fam Pract 2004; 21:575–581
- 24 Jennett PA, Laxdal OE, Hayton RC, et al. The effects of continuing medical education on family doctor performance in office practice: a randomized control study. Med Educ 1988; 22:139–145
- 25 Kottke TE, Brekke ML, Solberg LI, et al. A randomized trial to increase smoking intervention by physicians: doctors helping smokers, round I. JAMA 1989; 261:2101–2106
- 26 Lane DS, Messina CR, Grimson R. An educational approach to improving physician breast cancer screening practices and counseling skills. Patient Educ Couns 2001; 43:287–299
- 27 Lewis CE, Bursch B, Klau M, et al. Continuing medical education for AIDS: an organizational response. AIDS Educ Prev 1993; 5:263–271
- 28 Lindsay-McIntyre E, Wilson D, Best JA, et al. The impact of a continuing education package for smoking cessation on physicians' clinical behavior and patient smoking. Proc Annu Conf Res Med Educ 1987; 26:14–19
- 29 Maclure M, Dormuth C, Naumann T, et al. Influences of educational interventions and adverse news about calciumchannel blockers on first-line prescribing of antihypertensive drugs to elderly people in British Columbia. Lancet 1998; 352:943–948
- 30 Maiman LA, Becker MH, Liptak GS, et al. Improving pediatricians' compliance-enhancing practices: a randomized trial. Am J Dis Child 1988; 142:773–779
- 31 Mann KV, Lindsay EA, Putnam RW, et al. Increasing physician involvement In cholesterol-lowering practices: the role of knowledge, attitudes and perceptions. Adv Health Sci Educ Theory Pract 1997; 2:237–253
- 32 Margolis PA, Lannon CM, Stuart JM, et al. Practice based education to improve delivery systems for prevention in primary care: randomised trial. BMJ 2004; 328:388
- 33 McClellan WM, Millman L, Presley R, et al. Improved diabetes care by primary care physicians: results of a grouprandomized evaluation of the Medicare Health Care Quality Improvement Program (HCQIP). J Clin Epidemiol 2003; 56:1210-1217
- 34 Mehler PS, Krantz MJ, Lundgren RA, et al. Bridging the quality gap in diabetic hyperlipidemia: a practice-based intervention. Am J Med 2005; 118:1414
- 35 Moran JA, Kirk P, Kopelow M. Measuring the effectiveness of a pilot continuing medical education program. Can Fam Physician 1996; 42:272–276
- 36 Norris SL, Grothaus LC, Buchner DM, et al. Effectiveness of physician-based assessment and counseling for exercise in a staff model HMO. Prev Med 2000; 30:513–523
- 37 Ockene IS, Hebert JR, Ockene JK, et al. Effect of training and a structured office practice on physician-delivered nutrition counseling: the Worcester-Area Trial for Counseling in Hyperlipidemia (WATCH). Am J Prev Med 1996; 12:252–258
- 38 Ozer EM, Adams SH, Lustig JL, et al. Increasing the screening and counseling of adolescents for risky health behaviors: a primary care intervention. Pediatrics 2005; 115:960–968

- 39 Perera DR, LoGerfo JP, Shulenberger E, et al. Teaching sigmoidoscopy to primary care physicians: a controlled study of continuing medical education. J Fam Pract 1983; 16:785–788
- 40 Rabin DL. Adapting an effective primary care provider STD/HIV prevention training programme. AIDS Care 1998; 10(suppl):S75–S82
- 41 Rahme E, Choquette D, Beaulieu M, et al. Impact of a general practitioner educational intervention on osteoarthritis treatment in an elderly population. Am J Med 2005; 118:1262–1270
- 42 Ray WA, Stein CM, Byrd V, et al. Educational program for physicians to reduce use of non-steroidal anti-inflammatory drugs among community-dwelling elderly persons: a randomized controlled trial. Med Care 2001; 39:425–435
- 43 Rodney WM, Albers G. Flexible sigmoidoscopy: primary care outcomes after two types of continuing medical education. Am J Gastroenterol 1986; 81:133–137
- 44 Schectman JM, Schroth WS, Verme D, et al. Randomized controlled trial of education and feedback for implementation of guidelines for acute low back pain. J Gen Intern Med 2003; 18:773–780
- 45 Schroy PC, Heeren T, Bliss CM, et al. Implementation of on-site screening sigmoidoscopy positively influences utilization by primary care providers. Gastroenterology 1999; 117:304–311
- 46 Schwartzberg JG, Guttman R. Effect of training on physician attitudes and practices in home and community care of the elderly. Arch Fam Med 1997; 6:439–444
- 47 Soumerai SB, Avorn J. Predictors of physician prescribing change in an educational experiment to improve medication use. Med Care 1987; 25:210–221
- 48 Stein CM, Griffin MR, Taylor JA, et al. Educational program for nursing home physicians and staff to reduce use of non-steroidal anti-inflammatory drugs among nursing home residents: a randomized controlled trial. Med Care 2001; 39:436-445
- 49 Stross JK, Bole GG. Evaluation of an educational program for primary care practitioners, on the management of osteo-arthritis. Arthritis Rheum 1985; 28:108–111
- 50 Terry PB, Wang VL, Flynn BS, et al. A continuing medical education program in chronic obstructive pulmonary diseases: design and outcome. Am Rev Respir Dis 1981; 123:42–46
- 51 White CW, Albanese MA, Brown DD, et al. The effectiveness of continuing medical education in changing the behavior of physicians caring for patients with acute myocardial infarction: a controlled randomized trial. Ann Intern Med 1985; 102:686–692
- 52 Bjornson DC, Rector TS, Daniels CE, et al. Impact of a drug-use review program intervention on prescribing after publication of a randomized clinical trial. Am J Hosp Pharm 1990; 47:1541–1546
- 53 Brown R, Bratton SL, Cabana MD, et al. Physician asthma education program improves outcomes for children of lowincome families. Chest 2004; 126:369–374
- 54 Casebeer LL, Klapow JC, Centor RM, et al. An intervention to increase physicians' use of adherence-enhancing strategies in managing hypercholesterolemic patients. Acad Med 1999; 74:1334–1339
- 55 Evans CE, Haynes RB, Birkett NJ, et al. Does a mailed continuing education program improve physician performance? Results of a randomized trial in antihypertensive care. JAMA 1986; 255:501–504
- 56 Gullion DS, et al. Management of hypertension in private practice: a randomized controlled trial in continuing medical education. J Contin Educ Health Prof 1988; 8:239–255
- 57 Hagen BF, Armstrong-Esther C, Quail P, et al. Neuroleptic and benzodiazepine use in long-term care in urban and rural

- Alberta: characteristics and results of an education intervention to ensure appropriate use. Int Psychogeriatr 2005; 17:631-652
- 58 Kim CS, Kristopaitis RJ, Stone E, et al. Physician education and report cards: do they make the grade? Results from a randomized controlled trial. Am J Med 1999; 107:556–560
- 59 Lin EH, Katon WJ, Simon GE, et al. Achieving guidelines for the treatment of depression in primary care: is physician education enough? Med Care 1997; 35:831–842
- 60 Lin EH, Simon GE, Katzelnick DJ, et al. Does physician education on depression management improve treatment in primary care? J Gen Intern Med 2001; 16:614–619
- 61 Pazirandeh M. Does patient partnership in continuing medical education (CME) improve the outcome in osteoporosis management? J Contin Educ Health Prof 2002; 22:142–151
- 62 Pinto BM, Goldstein MG, DePue JD, et al. Acceptability and feasibility of physician-based activity counseling: the PAL project. Am J Prev Med 1998; 15:95–102
- 63 Schectman JM, Kanwal NK, Schroth WS, et al. The effect of an education and feedback intervention on group-model and network-model health maintenance organization physician prescribing behavior. Med Care 1995; 33:139–144
- 64 Schectman JM, Schroth WS, Elinsky EG, et al. The effect of education and drug samples on antihistamine prescribing costs in an HMO. HMO Pract 1996; 10:119–122
- 65 Sibley JC, Sackett DL, Neufeld V, et al. A randomized trial of continuing medical education. N Engl J Med 1982; 306:511–515
- 66 Socolar RR, Raines B, Chen-Mok M, et al. Intervention to improve physician documentation and knowledge of child sexual abuse: a randomized, controlled trial. Pediatrics 1998; 101:817–824
- 67 Solomon DH, Katz JN, La Tourette AM, et al. Multifaceted intervention to improve rheumatologists' management of glucocorticoid-induced osteoporosis: a randomized controlled trial. Arthritis Rheum 2004; 51:383–387
- 68 Stewart M, Marshall JN, Ostbye T, et al. Effectiveness of case-based on-line learning of evidence-based practice guidelines. Fam Med 2005; 37:131–138
- 69 Thom DH. Training physicians to increase patient trust. J Eval Clin Pract 2000; 6:245–253
- 70 Levinson W, Roter D. The effects of two continuing medical education programs on communication skills of practicing primary care physicians. J Gen Intern Med 1993; 8:318–324
- 71 Zuckerman IH, Weiss SR, McNally D, et al. Impact of an educational intervention for secondary prevention of myocardial infarction on Medicaid drug use and cost. Am J Manag Care 2004; 10:493–500
- 72 Maxwell JA, Sandlow LJ, Bashook PG. Effect of a medical care evaluation program on physician knowledge and performance. J Med Educ 1984; 59:33–38
- 73 Mukohara K, Schwartz MD. Electronic delivery of research summaries for academic generalist doctors: a randomised trial of an educational intervention. Med Educ 2005; 39:402–409
- 74 Schectman JM, Elinsky EG, Pawlson LG. Effect of education and feedback on thyroid function testing strategies of primary care clinicians. Arch Intern Med 1991; 151:2163–2166
- 75 Beaulieu MD, Rivard M, Hudon E, et al. Comparative trial of a short workshop designed to enhance appropriate use of screening tests by family physicians. Can Med Assoc J 2002; 167:1241–1246
- 76 Carney PA, Dietrich AJ, Freeman DH Jr, et al. A standardized-patient assessment of a continuing medical education program to improve physicians' cancer-control clinical skills. Acad Med 1995; 70:52–58
- 77 Fordis M, King JE, Ballantyne CM, et al. Comparison of the instructional efficacy of internet-based CME with live inter-

- active CME workshops: a randomized controlled trial. JAMA 2005; 294:1043–1051
- 78 Gifford DR, Holloway RG, Frankel MR, et al. Improving adherence to dementia guidelines through education and opinion leaders: a randomized, controlled trial. Ann Intern Med 1999; 131:237–246
- 79 Kronick J, Blake C, Munoz E, et al. Improving on-line skills and knowledge: a randomized trial of teaching rural physicians to use on-line medical information. Can Fam Physician 2003; 49:312–317
- 80 Ray WA, Schaffner W, Federspiel CF. Persistence of improvement in antibiotic prescribing in office practice. JAMA 1985; 253:1774–1776
- 81 Tziraki C, Graubard BI, Manley M, et al. Effect of training on adoption of cancer prevention nutrition-related activities by primary care practices: results of a randomized, controlled study. J Gen Intern Med 2000; 15:155–162
- 82 Allison JJ, Kiefe CI, Wall T, et al. Multicomponent Internet continuing medical education to promote chlamydia screening. Am J Prev Med 2005; 28:285–290
- 83 Cherkin D, Deyo RA, Berg AO, et al. Evaluation of a physician education intervention to improve primary care for low-back pain: I. Impact on physicians. Spine 1991; 16:1168–1172
- 84 Goldstein MK, Lavori P, Coleman R, et al. Improving adherence to guidelines for hypertension drug prescribing: cluster-randomized controlled trial of general versus patient-specific recommendations. Am J Manag Care 2005; 11:677–685
- 85 Greenberg LW, Jewett LS. The impact of two teaching techniques on physicians' knowledge and performance. J Med Educ 1985; 60:390–396
- 86 Heale J, Davis D, Norman G, et al. A randomized controlled trial assessing the impact of problem-based versus didactic teaching methods in CME. Proc Annu Conf Res Med Educ 1988; 27:72–77
- 87 Howe HL, Lehnherr M, Katterhagen JG. Effects of physician outreach programs on rural-urban differences in breast cancer management. J Rural Health 1997; 13:109–117
- 88 Kutcher SP, Lauria-Horner BA, MacLaren CM, et al. Evaluating the impact of an educational program on practice patterns of Canadian family physicians interested in depression treatment. Prim Care Companion J Clin Psychiatry 2002; 4:224–231
- 89 Labelle M, Beaulieu M, Renzi P, et al. Integrating clinical practice guidelines into daily practice: impact of an interactive workshop on drafting of a written action plan for asthma patients. J Contin Educ Health Prof 2004; 24:39–49
- 90 Lockyer JM, Fidler H, Hogan DB, et al. Dual-track CME: accuracy and outcome. Acad Med 2002; 77:S61–S63
- 91 Mazmanian PE, Daffron SR, Johnson RE, et al. Information about barriers to planned change: a randomized controlled trial involving continuing medical education lectures and commitment to change. Acad Med 1998; 73:882–886
- 92 McBride P, Underbakke G, Plane MB, et al. Improving prevention systems in primary care practices: the Health Education and Research Trial (HEART). J Fam Pract 2000; 49:115–125
- 93 Pimlott NJ, Hux JE, Wilson LM, et al. Educating physicians to reduce benzodiazepine use by elderly patients: a randomized controlled trial. Can Med Assoc J 2003; 168:835–839
- 94 Rosenthal MS, Lannon CM, Stuart JM, et al. A randomized trial of practice-based education to improve delivery systems for anticipatory guidance. Arch Pediatr Adolesc Med 2005; 159:456–463
- 95 Winickoff RN, Coltin KL, Morgan MM, et al. Improving physician performance through peer comparison feedback. Med Care 1984; 22:527–534
- 96 Costanza ME, Zapka JG, Harris DR, et al. Impact of a

- physician intervention program to increase breast cancer screening. Cancer Epidemiol Biomarkers Prev 1992; 1:581–589
- 97 Sharif I, Oruwariye T, Cohen R, et al. Effectiveness of clinician training in smoking cessation counseling. Arch Pediatr Adolesc Med 2002; 156:944–945
- 98 Dormuth CR, Maclure M, Bassett K, et al. Effect of periodic letters on evidence-based drug therapy on prescribing behaviour: a randomized trial. Can Med Assoc J 2004; 171: 1057–1061
- 99 Myers RE, Turner B, Weinberg D, et al. Impact of a physician-oriented intervention on follow-up in colorectal cancer screening. Prev Med 2004; 38:375–381
- 100 Andersen SM, Harthorn BH. Changing the psychiatric knowledge of primary care physicians: the effects of a brief intervention on clinical diagnosis and treatment. Gen Hosp Psychiatry 1990; 12:177–190
- 101 Bloomfield HE, Nelson DB, van Ryn M, et al. A trial of education, prompts, and opinion leaders to improve prescription of lipid modifying therapy by primary care physicians for patients with ischemic heart disease. Qual Saf Health Care 2005; 14:258–263
- 102 Frush K, Hohenhaus S, Luo X, et al. Evaluation of a Web-based education program on reducing medication dosing error: a multicenter, randomized controlled trial. Pediatr Emerg Care 2006; 22:62–70
- 103 Gerrity MS, Cole SA, Dietrich AJ, et al. Improving the recognition and management of depression: is there a role for physician education? J Fam Pract 1999; 48:949–957
- 104 Goldberg HI, Deyo RA, Taylor VM, et al. Can evidence change the rate of back surgery? A randomized trial of community-based education. Eff Clin Pract 2001; 4:95–104
- 105 Grady KE, Lemkau JP, Lee NR, et al. Enhancing mammography referral in primary care. Prev Med 1997; 26:791–800
- 106 Juzych NS, Banerjee M, Essenmacher L, et al. Improvements in antimicrobial prescribing for treatment of upper respiratory tract infections through provider education. J Gen Intern Med 2005; 20:901–905
- 107 Lane DS, Polednak AP, Burg MA. Effect of continuing medical education and cost reduction on physician compliance with mammography screening guidelines. J Fam Pract 1991; 33:359–368
- 108 Gerstein HC, Reddy SS, Dawson KG, et al. A controlled evaluation of a national continuing medical education programme designed to improve family physicians' implementation of diabetes-specific clinical practice guidelines. Diabet Med 1999; 16:964–969
- 109 Fick DM, Maclean JR, Rodriguez NA, et al. A randomized study to decrease the use of potentially inappropriate medications among community-dwelling older adults in a southeastern managed care organization. Am J Manag Care 2004; 10:761– 768
- 110 White M, Michaud G, Pachev G, et al. Randomized trial of problem-based versus didactic seminars for disseminating evidence-based guidelines on asthma management to primary care physicians. J Contin Educ Health Prof 2004; 24:237–243
- 111 Peterson L, Tremblay G, Ewigman B, et al. The parental daily diary: a sensitive measure of the process of change in a child maltreatment prevention program. Behav Modif 2002; 26:627– 647
- 112 Mazmanian PE, Davis DA. Continuing medical education and the physician as a learner: guide to the evidence. JAMA 2002; 288:1057–1060
- 113 Fox RD. Using theory and research to shape the practice of continuing professional development. J Contin Educ Health Prof 2000; 20:238–246
- 114 Fox V. Patient teaching: understanding the needs of the adult learner. AORN J 1986; 44:234–238, 240, 242

Continuing Medical Education Effect on Practice Performance

Dave Davis and Robert Galbraith Chest 2009;135; 42S-48S DOI 10.1378/chest.08-2517

This information is current as of April 9, 2009

Updated Information Updated Information and services, including

& Services high-resolution figures, can be found at:

http://www.chestjournal.org/content/135/3 suppl/42S.ful

l.html

References This article cites 116 articles, 35 of which can be

accessed free at:

http://www.chestjournal.org/content/135/3 suppl/42

S.full.html#ref-list-1

Citations This article has been cited by 1 HighWire-hosted

articles:

http://www.chestjournal.org/content/135/3_suppl/42

S.full.html#related-urls

Open Access Freely available online through CHEST open access

option

Permissions & Licensing Information about reproducing this article in parts

(figures, tables) or in its entirety can be found online at: http://www.chestjournal.org/site/misc/reprints.xhtml

Reprints Information about ordering reprints can be found online:

http://www.chestjournal.org/site/misc/reprints.xhtml

Email alerting service Receive free email alerts when new articles cit this

article. sign up in the box at the top right corner of the

online article.

Images in PowerPoint

format

Figures that appear in CHEST articles can be

downloaded for teaching purposes in PowerPoint slide

format. See any online article figure for directions.

